

# WRIA 26, LOWER COWLITZ WATERSHED RIPARIAN CONDITIONS

MAP - 11a

## Legend

	Primary Roads		Poor Riparian Conditions		Lower Cowlitz
	Secondary Roads		Unknown Riparian Conditions		Toutle
	Light Duty		Good Riparian Conditions		Coweeman


This map was compiled by Lewis County Geographic Information Services. The base map was developed by the Washington State Department of Natural Resources by scanning and digitizing United States Geologic Survey 1:24,000 quadrangle maps. Riparian Conditions were derived using a 30-meter buffer on landcover data from WDNK along Type 1, 2, and 3 DNR classified streams with anomalous presence (see GIS Based Evaluation of Salmon Habitat in the Pacific Northwest by Lunetta et. al 1997 for a complete description of the landcover dataset). Accuracy of the map has not been verified, and it should be used for locational purposes only. Any possible discrepancies should be brought to the attention of Lewis County Geographic Information Services.

Projection: Lambert Conformal Conic Datum: 1927 North American Datum  
U.S.G.S. State Plane Zone 5626 Washington South Zone 06/30/2000

**GOOD:**  
Stream segments designated as having a 30-meter riparian buffer containing >70% coniferous crown cover of only mid-to-late seral stage trees.

**UNKNOWN:**  
Stream segments that did not fall into the good or poor categories but included early seral stage cover where conditions vary widely from good to poor.

**POOR:**  
Stream segments designated as having a 30-meter riparian buffer of <10% coniferous canopy cover and/or non-forested lands (urban, ag., and rangeland).





Location of WRIA 26



# WRIA 26, UPPER COWLITZ WATERSHED



Location of WRIA 26



## RIPARIAN CONDITIONS MAP - 11b

### Legend

- |   |                             |   |                 |  |                 |
|---|-----------------------------|---|-----------------|--|-----------------|
|  | Poor Riparian Conditions    |  | Primary Roads   |  | Upper Cowlitz   |
|  | Unknown Riparian Conditions |  | Secondary Roads |  | Cispus          |
|  | Good Riparian Conditions    |  | Light Duty      |  | Mayfield-Tilton |
|   |                             |   |                 |  | Riffe           |

**GOOD:**  
Stream segments designated as having a 30-meter riparian buffer containing >70% coniferous crown cover of only mid- to late seral stage trees.

**UNKNOWN:**  
Stream segments that did not fall into the good or poor categories but included early seral stage cover where conditions vary widely from good to poor.

**POOR:**  
Stream segments designated as having a 30-meter riparian buffer of <10% coniferous canopy cover and/or non-forested lands (urban, ag., and rangeland).

This map was compiled by Lewis County Geographic Information Services. The base map was developed by the Washington State Department of Natural Resources by scanning and digitizing United States Geologic Survey 1:24,000 quadrangle maps. Riparian Conditions were derived using a 30-meter buffer on landcover data from WDN along Type 1, 2, and 3 DNR classified streams with anadromous presence (see GIS Based Evaluation of Salmon Habitat in the Pacific Northwest by Lunetta et. al 1997 for a complete description of the landcover dataset). Accuracy of the map has not been verified, and it should be used for locational purposes only. Any possible discrepancies should be brought to the attention of Lewis County Geographic Information Services.

Projection: Lambert Conformal Conic Datum: 1927 North American Datum  
U.S.G.S. State Plane Zone 5626 Washington South Zone 06/30/2000

